

February 6, 2008

Hole Photoproduction in Insulating Copper Oxide

O. P. Sushkov*

School of Physics, The University of New South Wales, Sydney 2052, Australia

Abstract

Basing on $t - J$ model we calculate the $\mathbf{k}$ -dependence of a single hole photoproduction probability for CuO_2 plane at zero doping. We also discuss the radiation of spin-waves which can substantially deform the shape of photoemission spectra.

PACS numbers: 75.50.Ee, 75.10.Jm,

Typeset using REVTeX

Recent photoemission measurements¹ for insulating Copper Oxide $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ give an unique possibility to determine single hole dispersion from experiment. However, to compare with data one needs to calculate the probability of the hole creation at a given momentum and energy transfer. The purpose of the present work is to calculate this probability in the framework of $t - J$ model and elucidate the related question with momentum dependence of quasihole residue (see Refs.^{2,3}). In the conclusion we also comment on the shape of photoemission spectra.

It is widely accepted that low energy dynamics of CuO_2 planes in Copper Oxides is described by $t - J$ model (see Ref.⁴ for a review). At zero doping which corresponds to half filling this model is equivalent to Heisenberg antiferromagnet with long range Néel order in the ground state. We are interested in the process of an external photon kicking out an electron from the plane and creating a hole. The properties of a single hole are well established. For its description at $t/J \leq 5$ one can use selfconsistent Born approximation⁵. This approximation is justified by the absence of a single loop correction to the spin-wave vertex⁶⁻⁹. In some sense it is analog of the well known Migdal theorem for electron-phonon interaction. Hole dispersion has minima at $\mathbf{k} = (\pm\pi/2, \pm\pi/2)$ and it is almost degenerate along the face of magnetic Brillouin zone $\gamma_{\mathbf{k}} = \frac{1}{2}(\cos k_x + \cos k_y) \geq 0$. The hole wave function $\psi_{\mathbf{k}\sigma}$ is characterized by quasimomentum $\mathbf{k}$ defined inside the magnetic Brillouin zone, and pseudo-spin $\sigma = \uparrow\downarrow$. The pseudo-spin denotes the sublattice at which the hole is centered, and it is different from the usual spin. It is convenient to write the hole wave function in the form

$$\psi_{\mathbf{k}\sigma} = h_{\mathbf{k}\sigma}^\dagger |0\rangle, \quad (1)$$

where $|0\rangle$ is the ground state of the Heisenberg model, and $h_{\mathbf{k}\sigma}^\dagger$ is the creation operator of the composite hole. Let us also denote by d_{ns} the annihilation operator of an electron at the site n and with usual spin $s = \uparrow\downarrow$. Let us fix the pseudo-spin σ in Eq. (1), for example $\sigma = \uparrow$. The quasiparticle residue of the composite hole is given by

$$\mathcal{Z}_{\mathbf{k}} \propto \left| \langle \psi_{\mathbf{k}\uparrow} | \sum_{n \in \text{up}} d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n} |0\rangle \right|^2. \quad (2)$$

We stress that the summation in this equation is restricted by one sublattice because the quasispin is fixed. This is the residue defined in the Refs.⁵⁻⁹, and due to the Bloch theorem the exact relation

$$\mathcal{Z}_{\mathbf{k}+\mathbf{Q}} = \mathcal{Z}_{\mathbf{k}} \quad (3)$$

is valid for translation at the inverse vector of the magnetic sublattice, ($\mathbf{Q} = (\pm\pi, \pm\pi)$).

When a photon kicks out an electron from the system it does not separate the sublattices, therefore the external perturbation is

$$\sqrt{\frac{2}{N}} \sum_n d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n}, \quad (4)$$

and the probability of the process is proportional to

$$A_{\mathbf{k}} = \left| \langle \psi_{\mathbf{k}\uparrow} | \sqrt{\frac{2}{N}} \sum_n d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n} | 0 \rangle \right|^2. \quad (5)$$

Here N is number of sites in the lattice. Normalization of the external perturbation is rather arbitrary. We choose it in such a way that $A_{\mathbf{k}} = 1$ for Ising case and $t = 0$. The value defined by Eq.(5) is different from the residue (2). For the calculation of $A_{\mathbf{k}}$ it is convenient to use $l = 1$ string variational ansatz for the hole wave function¹⁰

$$h_{\mathbf{k}\uparrow}^{(1)\dagger} = \sqrt{\frac{2}{N}} \sum_{n \in \uparrow} \left(\nu_{\mathbf{k}} d_{n\uparrow} + \mu_{\mathbf{k}} S_n^- \sum_{\delta} \left[(1+v) - (u+v) \gamma_{\mathbf{k}} e^{i\mathbf{k}\delta} \right] d_{n+\delta\downarrow} \right) e^{i\mathbf{k}\mathbf{r}_n}, \quad (6)$$

where $\sum_{\delta}$ denotes the summation over nearest neighbours, and the coefficients are given by

$$\begin{aligned} \nu_{\mathbf{k}} &= \frac{1}{2} \left[\frac{\Delta + 2S_{\mathbf{k}}}{X S_{\mathbf{k}}} \right]^{1/2}, \\ \mu_{\mathbf{k}} &= \frac{t}{[Y S_{\mathbf{k}} (\Delta + 2S_{\mathbf{k}})]^{1/2}}, \\ S_{\mathbf{k}} &= \left(\Delta^2/4 + 4t^2 \left[(1+y) - (x+y) \gamma_{\mathbf{k}}^2 \right] \right)^{1/2}. \end{aligned} \quad (7)$$

We set $J = 1$. The parameters $\Delta = 1.33$, $x = 0.56$, $y = 0.14$, $X = 0.8$, $Y = 0.72$, $u = 0.42$, $v = 0.12$ are some combinations of the Heisenberg model spin-spin correlators calculated in the Ref.¹⁰. The residue (2) is proportional to the coefficient $\nu_{\mathbf{k}}^2$ in the wave function (6). We would like to stress that the string representation of the hole wave function is dual to the usual spin-wave representation, but the physical matrix elements are certainly independent of the representation (see below).

Keeping in mind the relations $d_{n\uparrow}^\dagger d_{n\uparrow} = \frac{1}{2} + S_z$ and $d_{n\downarrow}^\dagger d_{n\uparrow} = S_n^-$ one can easily calculate the probability of photoemission (5) in the minimal string approximation

$$A_{\mathbf{k}}^{(1)} = \left[\nu_{\mathbf{k}} \left(\frac{1}{2} + \sigma \right) - 8\mu_{\mathbf{k}} (1-u) q_1 \gamma_{\mathbf{k}} \right]^2. \quad (8)$$

Here $\sigma = |\langle 0|S_z|0\rangle| = 0.3$ is average magnetization, and $q_1 = \frac{1}{2}\langle 0|S_n^+S_{n+\delta}^-|0\rangle = -0.08$ is the nearest neighbour spin-spin correlator for the Heisenberg model. Due to the $\gamma_{\mathbf{k}}$ term in (8) $A_{\mathbf{k}+\mathbf{Q}}^{(1)} \neq A_{\mathbf{k}}^{(1)}$! At zero hopping ($t = 0$) $A_{\mathbf{k}}^{(1)} = \frac{1}{2} + \sigma = 0.8$. It is smaller than 1 because of spin quantum fluctuations in the initial state. This value agrees very well with result of spin-wave calculation¹¹ and finite lattice computation¹².

The $l = 1$ string ansatz (6) is a good approximation for the hole wave function at $t \leq 1$. For $t > 1$ further dressing by $l > 1$ strings, or, in other words, multy spin-wave virtual excitations is important. There are many multy spin-wave components in the wave function of a dressed hole. The assumption that these components do not give a coherent contribution in $A_{\mathbf{k}}$, but only reduce the probability of $h_{\mathbf{k}\sigma}^{(1)}$ configuration is quite reasonable. This is exactly what happens for the hole spin-wave vertex⁹. Therefore

$$A_{\mathbf{k}} = Z_{\mathbf{k}} A_{\mathbf{k}}^{(1)}, \quad (9)$$

where $Z_{\mathbf{k}}$ is the weight of $h_{\mathbf{k}\sigma}^{(1)}$ in the exact $h_{\mathbf{k}\sigma}$. The values of $Z_{\mathbf{k}=(\pi/2,\pi/2)} \approx Z_{\gamma_{\mathbf{k}} \approx 0}$ calculated in the Ref.⁹ for different t are presented in the Table I. Note that the quasiparticle residue (2) equals to $\mathcal{Z}_{\mathbf{k}} = Z_{\mathbf{k}} \nu_{\mathbf{k}}^2 (1/2 + \sigma)$. Using this equality one can also find $Z_{\mathbf{k}}$ from the results of Refs.⁶⁻⁸ where quasiparticle residue $\mathcal{Z}_{\mathbf{k}}$ was calculated. In Table I we present the ratio $A_{\mathbf{Q}}/A_0 = A_{\mathbf{Q}}^{(1)}/A_0^{(1)}$ for different values of t . We see that $\mathbf{k}$ -dependence qualitatively agrees with experimental data¹. We assume that $t/J \geq 0$. It is interesting that at negative t the ratio $A_{\mathbf{Q}}/A_0$ is inverted. At $t \leq 1$ our result for $A_{\mathbf{Q}}/A_0$ reasonably agrees with that found by finite lattice computations^{12,13} and numerical spin-wave analysis². However at $t = 3$ the ratio $A_{\mathbf{Q}}/A_0$ found in Refs.^{12,13} for finite lattices is bigger than that from Table I. Possible reason of this this disagreement is a variational nature of $l = 1$ string calculation. Therefore below we discuss more regular way of calculation of $A_{\mathbf{k}}$.

Let us use not string, but spin-wave picture considering hopping term by perturbation theory⁵⁻⁸. The wave function of a bare hole (i.e. at $t = 0$) is of the form $\psi_{\mathbf{k}\uparrow}^b \approx \sqrt{\frac{2}{N}} \sum_{n \in \uparrow} d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n} |0\rangle$. External perturbation is given by Eq. (4), and we can introduce bare vertex of single hole production.

$$V_h^b = \langle \psi_{\mathbf{k}\uparrow}^b | \sqrt{\frac{2}{N}} \sum_n d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n} |0\rangle \approx 1. \quad (10)$$

Here we neglect the spin quantum fluctuations in the initial state. External perturbation (4) can also produce hole + spin-wave final state. Let us denote by $\alpha_{\mathbf{q}}^\dagger$ the creation operator

of the spin-wave with momentum $\mathbf{q}$ and projection of spin $S_z = -1$ (see Ref.¹⁴ for review). The vertex of hole + spin-wave creation equals

$$\begin{aligned} V_{h,sw}^b &= \langle 0 | \alpha_{\mathbf{q}} \left(\sqrt{\frac{2}{N}} \sum_{m \in \downarrow} d_{m\downarrow}^\dagger e^{-i(\mathbf{k}-\mathbf{q})\mathbf{r}_m} \right) \left(\sqrt{\frac{2}{N}} \sum_n d_{n\uparrow} e^{i\mathbf{k}\mathbf{r}_n} \right) | 0 \rangle = \\ &= \frac{2}{N} \langle 0 | \alpha_{\mathbf{q}} \sum_{m \in \downarrow} S_m^- e^{i\mathbf{q}\mathbf{r}_m} | 0 \rangle = \sqrt{\frac{2}{N}} v_{\mathbf{q}}, \end{aligned} \quad (11)$$

where $v_{\mathbf{q}}$ is Bogoliubov parameter diagonalizing spin-wave Hamiltonian: $u_{\mathbf{q}} = \sqrt{\frac{1}{\omega_{\mathbf{q}}} + \frac{1}{2}}$, $v_{\mathbf{q}} = -\text{sign}(\gamma_{\mathbf{q}}) \sqrt{\frac{1}{\omega_{\mathbf{q}}} - \frac{1}{2}}$, $\omega_{\mathbf{q}} = 2\sqrt{1 - \gamma_{\mathbf{q}}^2}$, (see Ref.¹⁴) We stress that (11) is a bare vertex. It corresponds to instant production of hole + spin wave, but not production of hole with subsequent decay into hole + spin-wave. Note that $V_{h,sw}^b \rightarrow \infty$ at $\mathbf{q} \rightarrow 0$. The reason is that perturbation (4) does not correspond any quasiparticle of the system, and therefore usual Goldstone theorem is not applicable. The vertices (10) and (11) are presented at Fig. 1. Cross corresponds to external perturbation, solid line - to the hole, and dashed line - to the spin-wave.

Bare hole-spin wave vertex is equal to $g_{\mathbf{p},\mathbf{q}} = \sqrt{\frac{2}{N}} 2f_0(\gamma_{\mathbf{p}} u_{\mathbf{q}} + \gamma_{\mathbf{k}} v_{\mathbf{q}})$, with $f_0 = 2t$ (see e.g. Refs.⁵⁻⁸). It corresponds to the decay of the hole with momentum $\mathbf{k}$ into a hole with momentum $\mathbf{p} = \mathbf{k} - \mathbf{q}$ and a spin-wave with momentum $\mathbf{q}$.

Now we can easily calculate the first correction to the production vertex (10). It is given by the diagram presented at Fig.2.

$$\begin{aligned} \delta V_h^{(1)}(\epsilon = \epsilon_{\mathbf{k}}, \mathbf{k}) &= \sqrt{\frac{2}{N}} \sum_{\mathbf{q}} \frac{v_{\mathbf{q}} g_{\mathbf{k}-\mathbf{q},\mathbf{q}}}{\epsilon_{\mathbf{k}} - \epsilon_{\mathbf{k}-\mathbf{q}} - \omega_{\mathbf{q}}} = \\ &= -\frac{8t}{N} \sum_{\mathbf{q}} \frac{v_{\mathbf{q}}(\gamma_{\mathbf{k}-\mathbf{q}} u_{\mathbf{q}} + \gamma_{\mathbf{k}} v_{\mathbf{q}})}{\omega_{\mathbf{q}}} = \frac{4t}{N} \gamma_{\mathbf{k}} \sum_{\mathbf{q}} \left(\frac{1}{\omega_{\mathbf{q}}} - \frac{1}{2} \right) = 0.4 \cdot t \cdot \gamma_{\mathbf{k}}. \end{aligned} \quad (12)$$

This gives

$$A_{\mathbf{k}} = \left(V_h^b + \delta V_h^{(1)} \right)^2 = (1 + 0.4 t \gamma_{\mathbf{k}})^2. \quad (13)$$

It is in a very good agreement with string variational result (8) which gives $A_{\mathbf{k}} = 0.8(1 + 0.37 t \gamma_{\mathbf{k}})^2$ at $t \rightarrow 0$.

To perform spin-wave calculations at large t one has to remember that one loop correction to the hole-spin-wave vertex is absent⁶⁻⁹ and therefore selfconsistent Born (SB) approximation⁵ is valid. It means that only modification in comparison with naive perturbation theory is that in the diagram Fig. 2 one has to use dressed hole Green function found

in selfconsistent Born approximation. We carried out simplified computation of this type with dressed hole Green function in the form $\mathcal{Z}/(\epsilon - \epsilon_{\mathbf{k}} + i0)$. This is equivalent to the replacement of bare hole-spin-wave coupling constant $f_0 = 2t$ with the effective one $f = \mathcal{Z} \cdot f_0$, see Ref.⁹. The results of this computation at $t \leq 4$ reasonably agree with variational answer (8), (9). One can certainly calculate $A_{\mathbf{k}}$ more accurately using hole Green function found in Refs.⁶⁻⁸. However, we would like to stress that for detailed analysis of photoemission spectra $A_{\mathbf{k}}$ is not enough.

As we already discussed an external photon can produce hole + spin-wave as well as a single hole. Therefore the $\epsilon, \mathbf{k}$ dependence of photoemission spectrum is given by the imaginary part of diagrams presented at Fig.3, with solid line be hole Green function in selfconsistent Born approximation G_{SB} . The Fig. 3 represents new Green function corresponding to the external perturbation (4). The diagram Fig.3a gives usual $Im G_{SB}(\epsilon, \mathbf{k})$. The diagram Fig. 3b takes into account already discussed difference of $A_{\mathbf{k}}$ from pure residue $\mathcal{Z}_{\mathbf{k}}$, but still this contribution is proportional to $Im G_{SB}$. The diagram Fig.3c gives nonresonant background caused by the emission of spin-waves. And finally the diagrams Fig3d,e represent interference between background and resonant production. This interference shifts maximum of photoemission spectrum from $\epsilon = \epsilon_{\mathbf{k}}$. Whether this shift could explain the experimental spectra¹ is still an open question.

Before conclusion we would like to note that the “electron” creation operator is actually Zhang and Rice singlet creation operator.¹⁵ Therefore formfactor of this singlet $F_{\mathbf{k}}$ should be also taken into account. Calculation of $F_{\mathbf{k}}$ is not a problem because the wave function of the singlet is now well known¹⁶.

In the present work we calculate single hole photoproduction probability $A_{\mathbf{k}}$ in frameworks of $t - J$ model. In agreement with experiment $A_{\mathbf{k}}$ drops down outside magnetic Brillouin zone. We also point out that shape of photoemission spectrum is different from $Im G_{SB}(\epsilon, \mathbf{k})$, and due to the interference the difference is not just a smooth background.

I am very grateful to D. Poilblanc and G. Sawatzky for attracting my attention to this problem. This work has been done during my stay at the Laboratoire de Physique Quantique, Universite Paul Sabatier. I am gratefully acknowledge the hospitality and financial support.

REFERENCES

- * Also at the Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia
- ¹ B. O. Wells et al, Phys. Rev. Lett. **74**, 964 (1995).
- ² A. Angelucci, S. Sorella, D. Poilblanc, Phys. Lett. A **198**, 145 (1995).
- ³ S. Sorella, preprint SISSA 160/95/CM/MB.
- ⁴ E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).
- ⁵ C. L. Kane, P. A. Lee, and N. Read, Phys. Rev. B **39**, 6880 (1989).
- ⁶ G. Martinez and P. Horsch, Phys. Rev. B **44**, 317 (1991).
- ⁷ Z. Liu and E. Manousakis, Phys. Rev. B **45**, 2425 (1992).
- ⁸ J. Igarashi and P. Fulde, Phys. Rev. B **45**, 12357 (1992).
- ⁹ O. P. Sushkov, Phys. Rev. B **49**, 1250 (1994).
- ¹⁰ O. P. Sushkov, Solid State Communications **83**, 303 (1992).
- ¹¹ A. G. Mal'shukov and G. D. Mahan, Phys. Rev. Lett. **68**, 2200 (1992).
- ¹² D. Poilblanc, T. Ziman, H. J. Schulz, E. Dagotto, Phys. Rev. B **47** 14267 (1993).
- ¹³ D. Poilblanc, H. J. Schulz, T. Ziman, Phys. Rev. B **47** 3268 (1993).
- ¹⁴ E. Manousakis, Rev. Mod. Phys. **63**, 1, (1991).
- ¹⁵ F. C. Zhang and T. M. Rice, Phys. Rev. B **37**, 3759 (1988).
- ¹⁶ V. I. Belinicher and A. L. Chernyshev, Phys. Rev. B **47** 390 (1993).

TABLES

TABLE I. The weight $Z_{\gamma_{\mathbf{k}} \approx 0}$ of $l = 1$ string configuration in the dressed hole wave function, and the ratio of a single hole creation probabilities $A_{\mathbf{k}}$ at $\mathbf{k} = \mathbf{Q} = (\pm\pi, \pm\pi)$ and $\mathbf{k} = 0$ for different values of t/J

t/J	0	0.2	0.5	1	2	3	4
Z	1	0.96	0.88	0.79	0.63	0.53	0.45
$A_{\mathbf{Q}}/A_0$	1	1.3	1.9	2.5	3.3	3.7	3.9

Figure captions,

Fig.1 Bare vertices of a single hole creation and hole + spin-wave creation. The cross corresponds to external perturbation, solid line - to the hole, and dashed line - to the spin-wave.

Fig.2 First order correction to the single hole creation amplitude.

Fig.3 The diagrams contributing into photoemission spectrum. Solid line corresponds to the hole Green function in selfconsistent Born approximation..



Fig. 1



Fig. 2




Fig. 3